

# Tutorial 4

February 13, 2020

## Question 1

An interviewer is conducting a phone survey. Each call has probability 0.4 to be answered.

- (a) What is the average number of calls that need to be placed before the first answer?

From last time,  $X \sim \text{Geometric}(p = 0.4)$ . The mean of the geometric distribution is:

$$\mathbf{E}(X) = \frac{1-p}{p}$$

With  $p = 0.4$ , the average number of (no-answer) calls that need to be placed before the first answer is:

$$\mathbf{E}(X) = \frac{1-0.4}{0.4} = \frac{0.6}{0.4} = 1.5$$

- (b) What is the probability that the interviewer will have to make 10 calls before getting 3 answered?

Let  $X$  represent the number of failures before the third answered call (success). Then

$$X \sim \text{NBinom}(r = 3, p = 0.4)$$

To make 10 calls before the third success means that a total of 11 calls are placed. Of these 11 calls, the first 10 calls consists of 2 successes and 8 failures (in any order), followed by a success.

Since  $X$  counts the number of failures, we seek  $\mathbf{P}(X = 8)$ .

$$\begin{aligned}\mathbf{P}(X = 8) &= \binom{x+r-1}{r-1} (1-p)^x p^r \\ &= \binom{8+3-1}{3-1} (1-0.4)^8 (0.4)^3 \\ &= \binom{10}{2} (0.6)^8 (0.4)^3 \\ &= 0.0484\end{aligned}$$

Note that we deviate slightly from the textbook in how we interpret the phrase “10 calls before 3 answered”. (Compare the above solution with example 3.37 on page 129)

- (c) What is the average number of non-answered calls before he or she gets 3 calls answered?

The mean of the negative binomial distribution is:

$$\mathbf{E}(X) = \frac{r(1-p)}{p}$$

With  $r = 3$  and  $p = 0.4$ , the average number of non-answered calls before getting 3 answered is:

$$\mathbf{E}(X) = \frac{3(1-0.4)}{0.4} = \frac{3(0.6)}{0.4} = 4.5$$

## Question 2

Aircraft arrive at a small airport according to a Poisson process at a rate of  $\alpha = 5$  per hour.

- (a) What is the probability that during the next hour there will be 6 arrivals?

Let  $X$  represent the number of arrivals in the next hour ( $t = 1$ ). Then  $\mu = \alpha t = 5 \cdot 1 = 5$  and  $X \sim \text{Poisson}(\mu = 5)$ . The probability we seek is  $\mathbf{P}(X = 6)$ .

$$\begin{aligned}\mathbf{P}(X = 6) &= \frac{e^{-\mu} \cdot \mu^x}{x!} \\ &= \frac{e^{-5} \cdot 5^6}{6!} \\ &= 0.1462\end{aligned}$$

- (b) What is the probability that during the next two hours there will be 7 arrivals?

Let  $Y$  represent the number of arrivals in the next two hours ( $t = 2$ ). Then  $\mu = \alpha t = 5 \cdot 2 = 10$  and  $Y \sim \text{Poisson}(\mu = 10)$ . The probability we seek is  $\mathbf{P}(X = 7)$ .

$$\begin{aligned}\mathbf{P}(X = 7) &= \frac{e^{-10} \cdot 10^7}{7!} \\ &= 0.0901\end{aligned}$$

- (c) What is the average number of arrivals during the next 3 hours?

Let  $W$  represent the number of arrivals in the next three hours ( $t = 3$ ). Then  $\mu = \alpha t = 5 \cdot 3 = 15$  and  $W \sim \text{Poisson}(\mu = 15)$ . The mean of a Poisson distribution with parameter  $\mu$  is  $\mu$ . Therefore the average number of arrivals during the next three hours is:

$$\mathbf{E}(W) = \mu = 15$$

### Question 3

Of the people passing through an airport metal detector, 0.5% activate it. Let  $X$  be the number among a randomly selected group of 500 who activate the detector.

- (a) What is the exact distribution of  $X$ ? What is the approximate distribution of  $X$ ?

The exact distribution of  $X$  is  $\text{Binom}(n = 500, p = 0.005)$ . As a rule of thumb, for  $n > 50$  and  $np < 5$ , we can approximate the binomial distribution with a Poisson distribution. Here,  $n = 500 > 50$  and  $np = 500 \cdot 0.005 = 2.5 < 5$ . The Poisson parameter  $\mu = np = 2.5$ . Then the approximate distribution of  $X$  is:

$$X \dot{\sim} \text{Poisson}(\mu = 2.5)$$

- (b) Compute  $\mathbf{P}(X = 5)$  (using the approximate distribution).

$$\begin{aligned}\mathbf{P}(X = 5) &\approx \frac{e^{-2.5} \cdot 2.5^5}{5!} \\ &= 0.0668\end{aligned}$$

- (c) Compute  $\mathbf{P}(X \leq 5)$  (using the approximate distribution).

$$\begin{aligned}\mathbf{P}(X \leq 5) &= \mathbf{P}(X = 0) + \mathbf{P}(X = 1) + \dots + \mathbf{P}(X = 5) \\ &\approx \sum_{x=0}^5 \frac{e^{-2.5} \cdot 2.5^x}{x!} \\ &= e^{-2.5} \sum_{x=0}^5 \frac{2.5^x}{x!} \\ &= 0.9580\end{aligned}$$